


Figure 6.1. Prevalence of obesity among adults aged 20 years and over: United States, 1997–June 2007




NOTES: Obesity is defined as a Body Mass Index (BMI) of 30 kg/m² or more. The measure is based on self-reported height and weight. The analyses excluded people with unknown height or weight (about 4% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Early Release” and Table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2007 NHIS. The estimate for 2007 was based on data collected from January through June. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For the period January through June 2007, 26.6% (95% confidence interval = 25.49–27.63%) of U.S. adults aged 20 years and over were obese which was not significantly different from the 2006 estimate of 26.4%.
- The prevalence of obesity among U.S. adults aged 20 years and over has generally increased over time from 19.4% in 1997 to 26.6% for the period January through June 2007.

Figure 6.2. Prevalence of obesity among adults aged 20 years and over, by age group and sex: United States, January–June 2007




NOTES: Obesity is defined as a Body Mass Index (BMI) of 30 kg/m² or more. The measure is based on self-reported height and weight. The analyses excluded 782 people (6.3%) with unknown height or weight.

DATA SOURCE: Based on data collected from January through June in the Sample Adult Core component of the 2007 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the prevalence of obesity was higher among adults aged 40–59 years (30.2%) than among adults aged 20–39 years (23.8%) and 60 years and over (25.0%).
- There was no significant difference in the prevalence of obesity between women and men for all three age groups.

Figure 6.3. Age-adjusted prevalence of obesity among adults aged 20 years and over, by sex and race/ethnicity: United States, January–June 2007



NOTES: Obesity is defined as a Body Mass Index (BMI) of 30 kg/m² or more. The measure is based on self-reported height and weight. The analyses excluded 782 people (6.3%) with unknown height or weight. Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 20–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: Based on data collected from January through June in the Sample Adult Core component of the 2007 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ Non-Hispanic black women were more likely than Hispanic and non-Hispanic white women to be obese.

■ Among the six sex-specific race/ethnicity groups, the age-adjusted prevalence of obesity was highest among non-Hispanic black women.

Data tables for Figures 6.1–6.3:

Data table for Figure 6.1. Prevalence of obesity among adults aged 20 years and over: United States, 1997–June 2007

Year	Percent (95% confidence interval)	
	Crude ¹	Age-adjusted ²
1997	19.4 (18.9-19.9)	19.5 (18.9-20.0)
1998	20.6 (20.1-21.1)	20.6 (20.0-21.1)
1999	21.5 (20.9-22.1)	21.5 (20.9-22.1)
2000	21.8 (21.2-22.4)	21.8 (21.2-22.3)
2001	23.0 (22.4-23.6)	22.9 (22.3-23.5)
2002	23.9 (23.3-24.6)	23.8 (23.2-24.5)
2003	23.7 (23.1-24.3)	23.5 (22.9-24.2)
2004	24.5 (23.9-25.1)	24.3 (23.8-25.0)
2005	25.4 (24.77-26.09)	25.3 (24.66-25.96)
2006	26.4 (25.62-27.09)	26.2 (25.44-26.90)
January–June 2007	26.6 (25.49-27.63)	26.5 (25.44-27.61)

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population and using seven age groups: 20–29 years, 30–39 years, 40–49 years, 50–59 years, 60–69 years, 70–79 years, and 80 years and over.

NOTES: Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Early Release” and Table III in the Appendix for more details.

DATA SOURCE: NHIS, 1997–June 2007. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for Figure 6.2. Prevalence of obesity among adults aged 20 years and over, by age group and sex: United States, January–June 2007

Age and sex	Percent	95% confidence interval
20–39 years		
Total	23.8	22.11-25.58
Male	25.3	22.78-27.76
Female	22.4	20.33-24.43
40–59 years		
Total	30.2	28.44-31.96
Male	30.3	27.93-32.72
Female	30.1	27.86-32.30
60 years and over		
Total	25.0	23.28-26.78
Male	24.0	21.54-26.55
Female	25.9	23.41-28.30
20 years and over: crude¹		
Total	26.6	25.49-27.63
Male	27.0	25.42-28.50
Female	26.2	24.82-27.52
20 years and over: age-adjusted²		
Total	26.5	25.44-27.61
Male	26.8	25.29-28.36
Female	26.1	24.78-27.49

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population and using seven age groups: 20–29 years, 30–39 years, 40–49 years, 50–59 years, 60–69 years, 70–79 years, and 80 years and over.

DATA SOURCE: National Health Interview Survey, January–June 2007. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for Figure 6.3. Age-adjusted prevalence of obesity among adults aged 20 years and over, by sex and race/ethnicity: United States, January–June 2007

Sex and race/ethnicity	Percent¹	95% confidence interval
Male		
Hispanic or Latino	28.0	23.84-32.20
Not Hispanic or Latino:		
White, single race	26.4	24.50-28.34
Black, single race	28.9	25.09-32.78
Female		
Hispanic or Latino	31.5	27.72-35.29
Not Hispanic or Latino:		
White, single race	23.4	21.82-25.02
Black, single race	40.2	37.07-43.28

¹Estimates for this Healthy People 2010 Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 20–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: National Health Interview Survey, January–June 2007. Data are based on household interviews of a sample of the civilian noninstitutionalized population.